

## DAFTAR PUSTAKA

- Abdul Mudalip, S. K., Abu Bakar, M. R., Jamal, P., Adam, F., Che Man, R., Sulaiman, S. Z., Mohd Arshad, Z. I., & Md Shaarani, S. (2018). Effects of Solvents on Polymorphism and Shape of Mefenamic Acid Crystals. *MATEC Web of Conferences*, 150, 0–5. <https://doi.org/10.1051/matecconf/201815002004>
- Abdullah, A., & Tahar, N. (2022). *Dissolution Improvement of Mefenamic Acid Through Cocrystallization with Aminobenzoic Acid Using Liquid-Assisted Grinding Method* Peningkatan Disolusi Asam Mefenamat Melalui Kokristalisasi dengan Para-Aminobenzoic Acid Menggunakan Metode Liquid - Assisted. 5(2), 46–53.
- BPOM RI. (2022). Peraturan Badan Pengawas Obat Dan Makanan Nomor 11 Tahun 2022 Tentang Tata Laksana Uji Bioekivalensi. *Bpom Ri*, 1–16.
- Budiman, A., Nurlatifah, E., & Amin, S. (2016). Enhancement of Solubility and Dissolution Rate of Glibenclamide by Cocrystal Approach with Solvent Drop Grinding Method. *International Journal of Current Pharmaceutical Review and Research*, 7(5), 248–250. [www.ijcpr.com](http://www.ijcpr.com)
- Depkes RI. (2014). *Farmakope Indonesia Edisi V*.
- Dixit, M., Kini, A. G., & Kulkarni, P. K. (2012). Enhancing the dissolution of polymorphs I and II of mefenamic acid by spray drying. *Turkish Journal of Pharmaceutical Sciences*, 9(1), 13–26.
- Fudholi, A., Muchtaridi, M., & Puspita, I. (2017). *Simvastatin-nicotinamide co-crystal : design , preparation and preliminary characterization*. 16(February), 297–303.
- Gozali, D., Bahti, H. H., & Soewandhi, S. N. (2012). Pembentukan Kokristal Antara Kalsium Atorvastatin dengan Isonikotinamid dan Karakterisasinya. *Jurnal Sains Materi Indonesia*, April, 103–110.
- Gozali, D., Putra, A. F. C., & Sopyan, I. (2014). Pengaruh Modifikasi Kristal Kalsium Atorvastatin Terhadap Laju Disolusi. *Bionatura-Jurnal Ilmu-Ilmu Hayati Dan Fisik*, 16(2), 83–88.
- Indra. (2014). *Upaya Penurunan Kompresibilitas Ramipril Melalui Teknik Rekayasa Kristal [Tesis]*. Bandung: ITB.
- Indra, I. (2017). Karakterisasi Padatan Hasil Proses Kokristalisasi Asam Mefenamat Menggunakan Metode Penguapan Pelarut. *Jurnal Kesehatan Bakti Tunas Husada: Jurnal Ilmu-Ilmu Keperawatan, Analisis Kesehatan Dan Farmasi*, 17(1), 21. <https://doi.org/10.36465/jkbth.v17i1.207>
- Isnaeni, N. L., Trisna Wulandari, W., & Alifiar, I. (2021). Pembuatan dan Karakterisasi Kokristal Kurkumin dengan Asam Askorbat Sebagai Koformer. *Prosiding Seminar Nasional Diseminasi Penelitian, September*,

- 122–129.
- Issa, M. G., & Ferraz, H. G. (2011). Intrinsic dissolution as a tool for evaluating drug solubility in accordance with the biopharmaceutics classification system. *Dissolution Technologies*, 18(3), 6–13. <https://doi.org/10.14227/DT180311P6>
- Jayasankar, A., Somwangthanaroj, A., Shao, Z. J., & Rodríguez-Hornedo, N. (2006). Cocrystal formation during cogrinding and storage is mediated by amorphous phase. *Pharmaceutical Research*, 23(10), 2381–2392. <https://doi.org/10.1007/s11095-006-9110-6>
- Khullar, R., Kumar, D., Seth, N., & Saini, S. (2012). Formulation and evaluation of mefenamic acid emulgel for topical delivery. *Saudi Pharmaceutical Journal*, 20(1), 63–67. <https://doi.org/10.1016/j.jpsp.2011.08.001>
- Kotak, U., Prajapati, V., Solanki, H., Jani, G., & Jha, P. (2015). Co-crystallization Technique - Its rationale and recent progress. *World Journal of Pharmacy and Pharmaceutical Sciences*, 4(04), 1484–1508. [www.wjpps.com/download/article/1427970181.pdf](http://www.wjpps.com/download/article/1427970181.pdf)
- Kumar, A., Singh, P., & Nanda, A. (2020). Hot stage microscopy and its applications in pharmaceutical characterization. *Applied Microscopy*, 50(1). <https://doi.org/10.1186/s42649-020-00032-9>
- Lee, S., Mellon, N., Shariff, A. M., & Leveque, J. (2017). *Aromatic Polyamines Covalent Triazine Polymer as Sorbent for CO<sub>2</sub> Adsorption*. 012173. <https://doi.org/10.1088/1757-899X/226/1/012173>
- Manuscript, A. (2015). *CrystEngComm*. <https://doi.org/10.1039/b000000x>
- Martindale, Sweetman, & C, S. (2007). Martindale: The Complete Drug Reference - 35th editionMartindale: The Complete Drug Reference - 35th edition. In London The Pharmaceutical Press, 2005. <https://doi.org/10.7748/ns2007.01.21.20.31.b571>
- Mirza, S., Miroshnyk, I., Heinamaki, J., & Yliruusi, J. (2008). Co - Crystals : an Emerging Approach for Enhancing Properties of Pharmaceutical Solids. *Dosis*, 24(02), 90–96. <http://www.pssrc.org/download-directory?task=document.viewdoc&id=6>
- Moffat, C. A., Osselton, D. M., & Widdop, B. (1996). Drugs and poisons. *Serials Librarian*, 27(2–3), 87–102. [https://doi.org/10.1300/J123v27n02\\_07](https://doi.org/10.1300/J123v27n02_07)
- Mustapa, D. R., Matematika, F., Ilmu, D. A. N., Alam, P., Studi, P., & Farmasi, E. (2012). *Universitas indonesia pengaruh suhu pembentukan*.
- Padrela, L., De Azevedo, E. G., & Velaga, S. P. (2012). Powder X-ray diffraction method for the quantification of cocrystals in the crystallization mixture. *Drug Development and Industrial Pharmacy*, 38(8), 923–929. <https://doi.org/10.3109/03639045.2011.633263>

- Panchagnula, R., Sundaramurthy, P., Pillai, O., Agrawal, S., & Raj, Y. A. (2004). Solid-State Characterization of Mefenamic Acid. *Journal of Pharmaceutical Sciences*, 93(4), 1019–1029. <https://doi.org/10.1002/jps.20008>
- Qiao, N., Li, M., Schlindwein, W., Malek, N., Davies, A., & Trappitt, G. (2011). Pharmaceutical cocrystals: An overview. *International Journal of Pharmaceutics*, 419(1–2), 1–11. <https://doi.org/10.1016/j.ijpharm.2011.07.037>
- Sanphui, P., Goud, N. R., Khandavilli, U. B. R., & Nangia, A. (2011). Fast dissolving curcumin cocrystals. *Crystal Growth and Design*, 11(9), 4135–4145. <https://doi.org/10.1021/cg200704s>
- Sari, Y. N., Zaini, E., & Ismed, F. (2019). Peningkatan Laju Disolusi Piperine dengan Pembentukan Multikomponen Kristal Menggunakan Asam Nikotinat. *Jurnal Sains Farmasi & Klinis*, 6(2), 180. <https://doi.org/10.25077/jsfk.6.2.180-185.2019>
- Savjani, K. T., Gajjar, A. K., & Savjani, J. K. (2012). Drug Solubility: Importance and Enhancement Techniques. *ISRN Pharmaceutics*, 2012(100 mL), 1–10. <https://doi.org/10.5402/2012/195727>
- Sekhon, B. (2009). Pharmaceutical co-crystals - a review. *ARS Pharmaceutica*, 50(Figure 1), 99–117.
- Settle, F. (1997). Handbook of instrumental techniques for analytical chemistry. In *Prentice Hall*. <https://doi.org/10.5860/choice.35-3308>
- Setyawan, D., & Isadiartuti, D. (2009). Karakterisasi Kompleks Inklusi Asam Mefenamat –  $\beta$ Siklodekstrin yang Dibuat dengan Metode Freeze Drying. *Jurnal Kefarmasian Indonesia* (pp. 1.1: 1-9).
- Thakuria, R., & Thakur, T. S. (2017). *Crystal Polymorphism in Pharmaceutical Science Author's personal copy. June*. <https://doi.org/10.1016/B978-0-12-409547-2.12570-3>
- Trask, A. V., & Jones, W. (2005). Crystal engineering of organic cocrystals by the solid-state grinding approach. *Topics in Current Chemistry*, 254, 41–70. <https://doi.org/10.1007/b100995>
- U.S Pharmacopoeia National Formulary* (Issue dm). (2021).
- Vishweshwar, P., A. McMahon, J., A Bis, J., & Michael J, Z. (2006). Pharmaceutical Co-Crystals. *Journal of Pharmaceutical Sciences*, 95(3), 499–516. <https://doi.org/10.1002/jps.20578>
- Weyna, D. R., Shattock, T., Vishweshwar, P., & Zaworotko, M. J. (2009). Synthesis and structural characterization of cocrystals and pharmaceutical cocrystals: Mechanochemistry vs slow evaporation from solution. *Crystal Growth and Design*, 9(2), 1106–1123. <https://doi.org/10.1021/cg800936d>
- Wichianphong, N., & Charoenchaitrakool, M. (2018). Application of Box –

Behnken design for processing of mefenamic acid – paracetamol cocrystals using gas anti-solvent ( GAS ) process. *Journal of CO<sub>2</sub> Utilization*, 26(February), 212–220. <https://doi.org/10.1016/j.jcou.2018.05.011>

Zaini, E., Halim, A., Soewandhi, S. N., & Dwi Setyawan. (2011). Peningkatan Laju Pelarutan Trimetoprim Melalui Metode Ko-Kristalisasi Dengan Nikotinamida. *Jurnal Farmasi Indonesia*, 5(July), 206–212.